

Oats Plant Probiotics

Oats Plant Probiotics Formulation by NTHRYS

Plant Probiotics Standardized by NTHRYS for Oats Cultivation addresses below parameters in order to assist farming community to achieve better profits in Oats Cultivation.

1. Increase in grain yield and quality.
2. It protects the crop from pests and diseases, such as Crown Rust, Powdery Mildew, and Stem Rust.
3. Enhancement in Nutrient Uptake and Metabolic Health.
4. Improving Plant Vigor and Grain Size.
5. Increases Leaf Size and Photosynthetic Efficiency.
6. Increases Chlorophyll Content.
7. Increases Soil Enzyme Activity and the Quantity and Biodiversity of Viable Microorganisms.
8. Enhancement in Rhizosphere Activity.
9. Solubilization of Phosphorus.
10. Solubilization of Potassium.
11. Increase Soil Fertility.

NTHRYS Research Team has formulated Oats Plant Probiotics Formulation using a battery of Microbial Consortium (> 20 strains) to address all the above Objectives.

NTHRYS Plant Probiotics should not be mistaken as Biofertilizers. They are next generation plant supporting microbiome to support respective farmers to gain good profits with their produce.

Application Process :

Soil Treatment Before crop initiation:

1. The soil is treated with **formulation**, before initiating Oats planting.
2. 3 Lit **offormulation**is mixed with 100 Kg of Farm Yard Manure (FYM) 1 to 2 days before sowing per acre.

Seed Treatment :

1. Oats seeds are treated with **formulation**.
2. The seeds are mixed with **formulation** for half an hour and dried under shade. (200 ml **offormulation** for 5kg of seeds).
3. 1 to 2 ml **offormulation** is dropped on the soil where seed is sown.

Transplanting: Not applicable for Oats as it is typically direct-seeded.

Disease Management in Oats:

1. Oats suffers from various pests and diseases like Aphids, Barley Yellow Dwarf Virus, and Oat Blast.
2. Spraying **formulation** on the plant as well as at the base would help prevent these diseases and pest infestations.

Probiotic Dosage at Various Growth Phases:

1. **Germination Phase:** 500 ml **offormulation**+ 150 lit water + 1.35 kg of Epsom salt one day before spraying per acre. This should be done after the Oats seeds have germinated.
2. **Vegetative Growth Phase:** 1 lit **offormulation**+ 200 lit water + 1.8 kg of Epsom salt one day before spraying per acre until plants are fully established.
3. **Flowering and Grain Formation Phase:** 2 lit **offormulation** added to 300 lit of water + 2.7 kg of Epsom salt one day before spraying per acre from the onset of flowering until grain maturity. **formulation** Spraying and **formulation**+ **FYM** mixture (3 Lit **offormulation** is mixed with 100 Kg of Farm Yard Manure (FYM) per acre) addition can be done once per every two weeks during active growth periods.

Issued by:

NTHRYS OPC PVT LTD

Ph: +91 - 7093184748

Web: www.nthrYS.com

Email: smo@nthrys.com

Last Updated: 20 May 2024 10:34 am