

Pansy Plant Probiotics

Pansy Plant Probiotics Formulationby NTHRYS

Plant Probiotics Standardized by NTHRYS for Pansy Cultivation addresses below parameters in order to assist farming community to achieve better profits in Pansy Cultivation.

- 1. Enhancement in bloom size, color vibrancy, and longevity.
- 2. It protects the plant from pests and diseases, such as Aphids, Powdery Mildew, and Pansy Downy Mildew.
- 3. Enhancement in Nutrient Uptake and Metabolic Health.
- 4. Improving Plant Vigor and Bloom Density.
- 5. Increases Leaf Size and Photosynthetic Efficiency.
- 6. Increases Chlorophyll Content.
- 7. Increases Soil Enzyme Activity and the Quantity and Biodiversity of Viable Microorganisms.
- 8. Enhancement in Rhizosphere Activity.
- 9. Solubilization of Phosphorus.
- 10. Solubilization of Potassium.
- 11. Increase Soil Fertility.

NTHRYS Research Team has formulated Pansy Plant Probiotics Formulationusing a battery of Microbial Consortium (> 20 strains) to address all the above Objectives.

NTHRYS Plant Probiotics should not be mistaken as Biofertilizers. They are next generation plant supporting microbiome to support respective farmers to gain good profits with their produce.

Application Process:

Soil Treatment Before crop initiation:



- 1. The soil is treated with **formulation**, before initiating Pansy planting.
- 2. 3 Lit of**formulation**is mixed with 100 Kg of Farm Yard Manure (FYM) 1 to 2 days before planting per acre.

Seed/Cutting Treatment:

- 1. Pansy seeds or cuttings are treated with**formulation**.
- 2. The seeds or cuttings are soaked in**formulation**for half an hour before planting. (Use a dilute solution of**formulation**for soaking).

Disease Management in Pansy:

- 1. Pansies suffer from various pests and diseases like Snails, Slugs, and Botrytis Blight.
- 2. Spraying **formulation** on the plant as well as at the base would help prevent these diseases and pest infestations.

Probiotic Dosage at Various Growth Phases:

- 1. **After Planting**: 500 ml of**formulation**+ 150 lit water + 1.35 kg of Epsom salt one day before spraying per acre. This should be done immediately after planting seeds or cuttings and before new growth begins.
- 2. **Vegetative Growth Phase**: 1 lit of**formulation**+ 200 lit water + 1.8 kg of Epsom salt one day before spraying per acre until plants are fully established.
- 3. **Pre-bloom and Bloom Phase:** 2 lit offormulation added to 300 lit of water + 2.7 kg of Epsom salt one day before spraying per acre. Repeat application during the pre-bloom and bloom stages to enhance bloom size, color vibrancy, and longevity. **formulation**Spraying and **formulation**+ **FYM** mixture (3 Lit offormulation mixed with 100 Kg of Farm Yard Manure (FYM) per acre) addition can be done once per every two weeks during active growth periods.

Issued by:

NTHRYS OPC PVT LTD

Ph: +91 - 7093184748 Web: www.nthrys.com Email: smo@nthrys.com

Last Updated: 20 May 2024 2:39 pm