

Saffron Plant Probiotics

Saffron Plant Probiotics Formulation by NTHRYS

Plant Probiotics Standardized by NTHRYS for Saffron Cultivation addresses below parameters in order to assist farming community to achieve better profits in Saffron Cultivation.

1. Increase in saffron stigma yield and quality.
2. It protects the crop from pests and diseases, such as Corm Rot, Saffron Rust, and Thrips.
3. Enhancement in Nutrient Uptake and Metabolic Health.
4. Improving Plant Vigor and Flower Production.
5. Increases Leaf Size and Photosynthetic Efficiency.
6. Increases Chlorophyll Content.
7. Increases Soil Enzyme Activity and the Quantity and Biodiversity of Viable Microorganisms.
8. Enhancement in Rhizosphere Activity.
9. Solubilization of Phosphorus.
10. Solubilization of Potassium.
11. Increase Soil Fertility.

NTHRYS Research Team has formulated Saffron Plant Probiotics Formulation using a battery of Microbial Consortium (> 20 strains) to address all the above Objectives.

NTHRYS Plant Probiotics should not be mistaken as Biofertilizers. They are next generation plant supporting microbiome to support respective farmers to gain good profits with their produce.

Application Process :

Soil Treatment Before crop initiation:

1. The soil is treated with **formulation**, before initiating Saffron planting.

2. 3 Lit of **formulation** is mixed with 100 Kg of Farm Yard Manure (FYM) 1 to 2 days before planting per acre.

Corm Treatment :

1. Saffron corms are treated with **formulation**.
2. The corms are soaked in **formulation** for half an hour before planting. (Use a dilute solution of **formulation** for soaking).

Disease Management in Saffron:

1. Saffron suffers from various pests and diseases like Saffron Crocus Chlorotic Ringspot Virus, Bulb Rot, and Nematodes.
2. Spraying **formulation** on the plant as well as at the base would help prevent these diseases and pest infestations.

Probiotic Dosage at Various Growth Phases:

1. **After Planting:** 500 ml of **formulation** + 150 lit water + 1.35 kg of Epsom salt one day before spraying per acre. This should be done immediately after planting the corms and before the growth of new shoots begins.
2. **Vegetative Growth Phase:** 1 lit of **formulation** + 200 lit water + 1.8 kg of Epsom salt one day before spraying per acre until plants are fully established.
3. **Pre-harvest Phase:** 2 lit of **formulation** added to 300 lit of water + 2.7 kg of Epsom salt one day before spraying per acre a few weeks before the flowering period to enhance stigma development and quality. **formulation** Spraying and **formulation** + **FYM** mixture (3 Lit of **formulation** is mixed with 100 Kg of Farm Yard Manure (FYM) per acre) addition can be done once per every two weeks during active growth periods.

Issued by:

NTHRYS OPC PVT LTD

Ph: +91 - 7093184748

Web: www.nthrys.com

Email: smo@nthrys.com

Last Updated: 20 May 2024 2:35 pm