

Snake Gourd Plant Probiotics

Snake Gourd Plant Probiotics Formulation by NTHRYS

Plant Probiotics Standardized by NTHRYS for Snake Gourd Cultivation addresses below parameters in order to assist farming community to achieve better profits in Snake Gourd Cultivation.

1. Increase in fruit yield and quality.
2. It protects the crop from pests and diseases, such as Powdery Mildew, Mosaic Virus, and Fruit Fly.
3. Enhancement in Nutrient Uptake and Metabolic Health.
4. Improving Plant Vigor and Fruit Size.
5. Increases Leaf Size and Photosynthetic Efficiency.
6. Increases Chlorophyll Content.
7. Increases Soil Enzyme Activity and the Quantity and Biodiversity of Viable Microorganisms.
8. Enhancement in Rhizosphere Activity.
9. Solubilization of Phosphorus.
10. Solubilization of Potassium.
11. Increase Soil Fertility.

NTHRYS Research Team has formulated Snake Gourd Plant Probiotics Formulation using a battery of Microbial Consortium (> 20 strains) to address all the above Objectives.

NTHRYS Plant Probiotics should not be mistaken as Biofertilizers. They are next generation plant supporting microbiome to support respective farmers to gain good profits with their produce.

Application Process :

Soil Treatment Before crop initiation:

1. The soil is treated with **formulation**, before initiating Snake Gourd planting.
2. 3 Lit **offormulationis** mixed with 100 Kg of Farm Yard Manure (FYM) 1 to 2 days before planting per acre.

Seed Treatment :

1. Snake Gourd seeds are treated with **formulation**.
2. The seeds are mixed with **formulation** for half an hour and dried under shade. (200 ml **offormulation** for 5kg of seeds).
3. 1 to 2 ml **offormulationis** dropped on the soil where seed is sown.

Sowing: Directly in the field after seed treatment.

Disease Management in Snake Gourd:

1. Snake Gourd suffers from various pests and diseases like Cucumber Beetles, Anthracnose, and Downy Mildew.
2. Spraying **formulation** on the plant as well as at the base and on the leaves would help prevent these diseases and pest infestations.

Probiotic Dosage at Various Growth Phases:

1. **Germination Phase:** 500 ml **offormulation**+ 150 lit water + 1.35 kg of Epsom salt one day before spraying per acre. This should be done after the Snake Gourd seeds have germinated.
2. **Vegetative Growth Phase:** 1 lit **offormulation**+ 200 lit water + 1.8 kg of Epsom salt one day before spraying per acre until plants are fully established.
3. **Pre-harvest Phase:** 2 lit **offormulation** added to 300 lit of water + 2.7 kg of Epsom salt one day before spraying per acre a few weeks before the first harvest to enhance fruit development and quality. **formulation** Spraying and **formulation**+ **FYM** mixture (3 Lit **offormulationis** mixed with 100 Kg of Farm Yard Manure (FYM) per acre) addition can be done once per every two weeks during active growth periods.

Issued by:

NTHRYS OPC PVT LTD

Ph: +91 - 7093184748

Web: www.nthryst.com

Email: smo@nthryst.com

Last Updated: 10 May 2024 5:20 pm