

## Sorghum Plant Probiotics

# Sorghum Plant Probiotics Formulation by NTHRYS

Plant Probiotics Standardized by NTHRYS for Sorghum Cultivation addresses below parameters in order to assist farming community to achieve better profits in Sorghum Cultivation.

1. Increase in grain yield and quality.
2. It protects the crop from pests and diseases, such as Sorghum Midge, Anthracnose, and Leaf Blight.
3. Enhancement in Nutrient Uptake and Metabolic Health.
4. Improving Plant Vigor and Grain Size.
5. Increases Leaf Size and Photosynthetic Efficiency.
6. Increases Chlorophyll Content.
7. Increases Soil Enzyme Activity and the Quantity and Biodiversity of Viable Microorganisms.
8. Enhancement in Rhizosphere Activity.
9. Solubilization of Phosphorus.
10. Solubilization of Potassium.
11. Increase Soil Fertility.

NTHRYS Research Team has formulated Sorghum Plant Probiotics Formulation using a battery of Microbial Consortium (> 20 strains) to address all the above Objectives.

NTHRYS Plant Probiotics should not be mistaken as Biofertilizers. They are next generation plant supporting microbiome to support respective farmers to gain good profits with their produce.

### **Application Process :**

#### **Soil Treatment Before crop initiation:**

1. The soil is treated with **formulation**, before initiating Sorghum planting.
2. 3 Lit **offormulationis** mixed with 100 Kg of Farm Yard Manure (FYM) 1 to 2 days before planting per acre.

**Seed Treatment :**

1. Sorghum seeds are treated with **formulation**.
2. The seeds are mixed with **formulation** for half an hour and dried under shade. ( 200 ml **offormulation** for 5kg of seeds ).
3. 1 to 2 ml **offormulationis** dropped on the soil where seed is sown.

**Transplanting:** Not applicable for Sorghum as it is typically direct-seeded.

**Disease Management in Sorghum:**

1. Sorghum suffers from various pests and diseases like Sorghum Head Smut, Striga, and Sorghum Rust.
2. Spraying **formulation** on the plant as well as at the base would help prevent these diseases and pest infestations.

**Probiotic Dosage at Various Growth Phases:**

1. **Germination Phase:** 500 ml **offormulation**+ 150 lit water + 1.35 kg of Epsom salt one day before spraying per acre. This should be done after the Sorghum seeds have germinated.
2. **Vegetative Growth Phase:** 1 lit **offormulation**+ 200 lit water + 1.8 kg of Epsom salt one day before spraying per acre until plants are fully established.
3. **Pre-harvest Phase:** 2 lit **offormulation** added to 300 lit of water + 2.7 kg of Epsom salt one day before spraying per acre a few weeks before harvesting to enhance grain development and quality. **formulation** Spraying and **formulation**+ **FYM** mixture ( 3 Lit **offormulationis** mixed with 100 Kg of Farm Yard Manure (FYM) per acre) addition can be done once per every two weeks during active growth periods.

---

*Issued by:*

**NTHRYS OPC PVT LTD**

Ph: +91 - 7093184748

Web: [www.nthryst.com](http://www.nthryst.com)

Email: [smo@nthryst.com](mailto:smo@nthryst.com)

Last Updated: 11 May 2024 1:45 am