



## Tamarind Plant Probiotics

# Tamarind Plant Probiotics Formulation by NTHRYS

Plant Probiotics Standardized by NTHRYS for Tamarind Cultivation addresses below parameters in order to assist farming community to achieve better profits in Tamarind Cultivation.

1. Increase in fruit yield and quality.
2. It protects the crop from pests and diseases, such as Tamarind Seed Borer, Powdery Mildew, and Fusarium Wilt.
3. Enhancement in Nutrient Uptake and Metabolic Health.
4. Improving Plant Vigor and Fruit Size.
5. Increases Leaf Size and Photosynthetic Efficiency.
6. Increases Chlorophyll Content.
7. Increases Soil Enzyme Activity and the Quantity and Biodiversity of Viable Microorganisms.
8. Enhancement in Rhizosphere Activity.
9. Solubilization of Phosphorus.
10. Solubilization of Potassium.
11. Increase Soil Fertility.

NTHRYS Research Team has formulated Tamarind Plant Probiotics Formulation using a battery of Microbial Consortium (> 20 strains) to address all the above Objectives.

NTHRYS Plant Probiotics should not be mistaken as Biofertilizers. They are next generation plant supporting microbiome to support respective farmers to gain good profits with their produce.

### **Application Process :**

### **Soil Treatment Before crop initiation:**

1. The soil is treated with **formulation**, before initiating Tamarind planting.
2. 3 Lit **offormulationis** mixed with 100 Kg of Farm Yard Manure (FYM) 1 to 2 days before planting per acre.

#### **Seedling/Cutting Treatment :**

1. Tamarind seedlings or cuttings are treated with **formulation**.
2. The seedlings or cuttings are soaked in **formulation** for half an hour before planting. (Use a dilute solution **offormulation** for soaking).

#### **Disease Management in Tamarind:**

1. Tamarind suffers from various pests and diseases like Leaf Spot, Tamarindus Fruit Borer, and Root Rot.
2. Spraying **formulation** on the plant as well as at the base and on the leaves would help prevent these diseases and pest infestations.

#### **Probiotic Dosage at Various Growth Phases:**

1. **After Planting:** 500 ml **offormulation**+ 150 lit water + 1.35 kg of Epsom salt one day before spraying per acre. This should be done immediately after planting the seedlings or cuttings and before new growth begins.
2. **Vegetative Growth Phase:** 1 lit **offormulation**+ 200 lit water + 1.8 kg of Epsom salt one day before spraying per acre until plants are fully established.
3. **Pre-harvest Phase:** 2 lit **offormulation** added to 300 lit of water + 2.7 kg of Epsom salt one day before spraying per acre a few weeks before the first harvest to enhance fruit development and quality. **formulation** Spraying and **formulation**+ **FYM** mixture ( 3 Lit **offormulationis** mixed with 100 Kg of Farm Yard Manure (FYM) per acre) addition can be done once per every two weeks during active growth periods.

---

*Issued by:*

**NTHRYS OPC PVT LTD**

Ph: +91 - 7093184748

Web: [www.nthryst.com](http://www.nthryst.com)

Email: [smo@nthryst.com](mailto:smo@nthryst.com)

Last Updated: 09 May 2024 9:26 am